

EXTRAORDINARILY HOT X-RAY EMISSION
FROM THE O9 EMISSION LINE STAR HD 119682CARA E. RAKOWSKI¹, N. S. SCHULZ², S. J. WOLK¹, AND PAOLA TESTA²*Draft version February 5, 2008*

ABSTRACT

We present new optical and X-ray observations to show that the X-ray source 1WGA J1346.5–6255 previously associated with the SNR G309.2-0.6 can be unequivocally identified with the emission line star HD 119682 located in the foreground open cluster NGC 5281. Images from *Chandra* in the X-ray band as well as from *Magellan* in the narrow optical H α band show a coincidence of the source positions within 0.5". The X-ray source appears extremely hot for an OB-star identified as of O9.7e type. *XMM-Newton* spectra show plasma temperatures of 1 keV and > 8 keV with an X-ray luminosity of $6.2 \pm 0.1 \times 10^{32}$ erg s⁻¹. The optical and X-ray properties are very reminiscent of the prototype emission line star γ Cas. We discuss the ramifications of this similarity with respect to very early type emission line stars as a new class of hard X-ray sources.

Subject headings: stars:emission-line, Be — stars:individual(HD 119682) — X-rays:stars

1. INTRODUCTION

X-ray emission from early type stars has many origins and much progress has been made, particularly in recent years, to understand some of the underlying mechanisms. Since the discovery with the *Einstein Observatory* (Harnden & et al. 1979; Seward et al. 1979) it has been established that O-stars and most early-type B-stars emit X-rays of some form (Pallavicini et al. 1981; Chlebowski et al. 1989; Berghoefer & Schmitt 1994; Cassinelli et al. 1994). Observations with *ROSAT* established a fairly tight relationship between the X-ray and bolometric luminosity with $L_X/L_{bol} \sim 10^{-7}$ (Berghoefer et al. 1996). OB-stars that follow this relation are thought to generate X-rays from a distribution of shocks in a line driven wind (Lucy & White 1980; Lucy 1982; Owocki et al. 1988). We have a relatively fair understanding how the shocks lead to X-ray emission and for involved wind velocities one can derive X-ray emissivities from plasma temperatures of up to 25 MK. However, once it comes to processes that lead to these shocks, models differ quite substantially (Owocki et al. 1988; Feldmeier 1995; Howk et al. 2000; Feldmeier & Shlosman 2002).

Soon after the launch of *Chandra* and *XMM-Newton* it became clear that the origins of X-rays in hot stars is much more complex and puzzling. Some early observations confirmed the picture where X-rays arise from shock fragments embedded within about 10 stellar radii of the stellar wind (Kahn et al. 2001; Waldron & Cassinelli 2001; Cassinelli et al. 2001; Miller et al. 2002), but others showed unusually high plasma temperatures and X-ray line properties which seemed to be more related to forms of magnetic activity (Schulz et al. 2000; Waldron & Cassinelli 2001; Cohen et al. 2003; Schulz et al. 2003; Gagné et al. 2005; Schulz et al. 2006). Hard X-rays from hot stars have now been observed

from various circumstellar environments. Enhanced X-ray emission from magnetically confined winds has been observed in young OB-stars in and near the Orion Trapezium (Schulz et al. 2003; Gagné et al. 2005; Schulz et al. 2006) and also seems the likely process for τ Sco (Cohen et al. 2003; Donati & et al. 2006). Candidates for X-ray enhancement through magnetic fields or colliding winds may be located in other young embedded clusters (Wolk et al. 2002; Rho et al. 2004; Townsley et al. 2003). Hard X-rays from colliding wind binaries have been observed in several exotic O-star binaries (Sana et al. 2004; Corcoran 2005; Pollock et al. 2005). Binary interactions can also lead to powerful hard X-ray outbursts when two magnetically active stars interact as has been recently proposed for the massive spectroscopic binary in θ^2 Ori A (Schulz et al. 2006).

A third category has recently been identified as a possible source of intrinsic hard X-rays: hot emission line stars. The prototype of such a source is γ Cas, which before *Chandra* was known as a hard X-ray source but was considered likely to be a binary system of a B0.5e star orbited by a degenerate companion, probably a white dwarf (Kubo et al. 1998; Owens et al. 1999). Smith et al. (2004) recently reported on high resolution spectra obtained with *Chandra* which indicated quite a number of peculiar X-ray properties. The spectra featured various emission line spectral components from hot collisional plasmas which range from 100 eV to 12 keV as well as cold fluorescence. In their analysis Smith et al. (2004) conclude that the X-ray emission is likely to originate close to the Be star and its disk.

In this paper we report on X-ray observations of another early-type star with bright, hot X-ray emission, the O9e emission line star HD 119682, located in the open cluster NGC 5281. The open cluster itself is fairly well-studied. Photometry and proper motion of NGC 5281 were investigated by Sanner et al. (2001), who found an age for the cluster of 45 ± 10 Myr and a distance of 1580 ± 150 pc from fits to the color-magnitude diagram after statistical field star subtraction. They find a membership probability $P = 0.78$ for HD 119682 based on the proper motion. Levenhagen & Leister (2004) spec-

Electronic address: crakowski@cfa.harvard.edu

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street Cambridge, MA 02138² Massachusetts Institute of Technology, Center for Space Research, 70 Vassar Street, Cambridge, MA 02139

troscopically determined the fundamental parameters for HD 119682, an effective temperature of 31910 ± 550 K, an age of 4 ± 1 Myr, and a luminosity $\log(L/L_{\odot}) = 4.64 \pm 0.1$ implying a distance modulus consistent with NGC 5281. Given updated determinations of O star effective temperatures which account for line-blanketing (Martins et al. 2002), this effective temperature categorizes HD 119682 as an O9.7 star, but it is certainly on the borderline between O and B, with Levenhagen & Leister (2006) recently interpreting their measurements as a B0Ve star. The optical spectra from Levenhagen & Leister (2004) do not show any evidence for binarity.

The unequivocal identification of the X-ray source 1WGA J1346.5–6255 (White et al. 1994b,a) with HD 119682 had previously been thwarted by the lack of sufficient spatial resolution in the X-ray band to eliminate source confusion in the center of NGC 5281 as well as the proximity on the sky of a young supernova remnant (SNR) G309.2–0.6 for which 1WGA J1346.5–6255 could potentially have been a pulsar-wind-nebula (PWN). Herein we present new *Chandra* X-ray and *Magellan* H α observations with the spatial resolution to unambiguously establish the correspondence between 1WGA J1346.5–6255 and HD 119682 and discuss the deep X-ray spectrum of HD 119682 taken with *XMM-Newton* that even at CCD spectral resolution reveals emission lines from highly ionized Fe indicative of a hot thermal plasma.

2. OBSERVATIONS

To obtain an accurate and more finely constrained X-ray source position we used an archival *Chandra* observation of the field. On 2004 December 26, NGC 5281 was observed for 15 ks by the backside illuminated chip (S3) of the Advanced CCD Imaging Spectrometer of *Chandra* (OBSID:4554). The data were reduced and preliminarily analyzed using the pipeline developed for the *Chandra* star formation archive ANCHORS (Spitzbart et al. 2005). With 1496 net photons from 1WGA J1346.5–6255, the *Chandra* spectrum suffers mildly from pile-up ($\sim 12\%$). In addition to the known bright X-ray source, 45 other point sources (19 with more than 20 net counts each) were detected, a number of which are coincident with other members of the open cluster. This field also contains a large portion of the supernova remnant (SNR) G309.2–0.6, believed to lie a few parsecs behind NGC 5281. The radio source ATCA J134649-625235, identified by Gaensler et al. (1998) as a candidate pulsar-wind-nebula (PWN) associated with this SNR is detected as an X-ray source, with 23 counts $3'$ away from HD 119682.

For the purposes of accurate astrometry and to reduce source confusion a 5 s narrow-band H α image of the central $2.3'$ of NGC 5281 was taken with the MagIC CCD camera on the 6.5 m Clay Telescope of the *Magellan* Observatory at Las Campanas, Chile on 2004 February 17 UT. The excellent $<0.45''$ seeing and $0.069''$ pixel scale allowed us to register the field relative to the Guide Star Catalogue-II to an accuracy of $<0.75''$ with 30 matched stars using WCSTools (Mink 2002). The resulting coordinates for HD 119682 were $13^{\text{h}}46^{\text{m}}32^{\text{s}}.56, -62^{\circ}55'24''.06$ (J2000). We can also conservatively rule out any bright optical companion further than $0.75''$ away simply from the lack of any non-instrumental distortions in the profile




FIG. 1.— H α image of open cluster NGC 5281 with *Chandra* X-ray contours overlaid. The H α image was taken with the MagIC camera at *Magellan* to better determine the positions of the bright stars. The contours shown are 1, 5, and 50 counts after adaptive smoothing. 1WGA J1346.5–6255 is clearly identified as the O9 emission line star HD 119682. (The radio source ATCA J134649-625235 and its *Chandra* X-ray counterpart lie $3'$ to the northwest of this image.)

of HD 119682 (~ 0.006 pc at the distance to NGC 5281).

A 39.9 ks *XMM-Newton* observation of SNR G309.2–0.6 was taken on 2001 August 28, OBSID:0087940201 for which NGC 5281 was in the field, $6.5'$ off-axis. Three significant background flares occurred during the observation affecting EPIC-PN slightly more than the MOS instruments, resulting in filtered exposure times of 21.9 ks and 27.0 ks for PN and MOS respectively. The events were filtered using the standard criteria and *XMM-SAS* tools. For spectra from the MOS instruments, patterns between 0 and 12 were kept. For the PN two spectra were extracted, one with only the single pixel events (pattern=0) and one for the multiple pixel events (patterns 1 to 4). The single pixel response is slightly better known and we can compare the single and multiple event spectra for signs of pile-up. Although extremely bright, we find that 1WGA J1346.5–6255 is not piled-up in any of the *XMM-Newton* instruments. A $30''$ radius region centered at the *Chandra* source position was used to extract the events from 1WGA J1346.5–6255. With 4068, 2677, 2696, and 2880 counts in EPIC PN single, EPIC PN multiple, MOS1, and MOS2, respectively, background subtraction is essentially negligible for 1WGA J1346.5–6255, reducing the count-rates by less than 5%.

3. X-RAY PROPERTIES OF HD 119682

ASCA observations showed that 1WGA J1346.5–6255 had a low column density more consistent with the distance to the open cluster NGC 5281 than SNR G309.2–0.6 (Rakowski et al. 2001). The *Chandra* ACIS-S observations show 1WGA J1346.5–6255 to be an unresolved source coincident with HD 119682 (Figure 1). The flux has remained relatively constant (within 30%) across the *ASCA*, *XMM-Newton*, and *Chandra* observations in 1999 April, 2001 August, and 2004 December. No periodicities were found from the power spectra of any of the light-curves of 1WGA J1346.5–6255. However the

TABLE 1
THERMAL (RAYMOND-SMITH) MODEL PARAMETERS

Parameter	Value	
	1 kT	2 kT
N_H (10^{21} atoms cm^{-2})	1.8 ± 0.1	1.9 ± 0.1
abundance ($\times$ solar)	0.39 ± 0.13	0.40 ± 0.17
$kT(\text{soft})(\text{keV})$	...	1.07 ± 0.16
flux(10^{-12} ergs cm^{-2} s^{-1}) ^a	...	0.08 ± 0.04
$kT(\text{hot})(\text{keV})$	$8.1^{+0.9}_{-0.8}$	$10.4^{+2.3}_{-1.6}$
flux(10^{-12} ergs cm^{-2} s^{-1}) ^a	1.81 ± 0.01	1.74 ± 0.03
$\chi^2(\text{d.o.f.})$	450.56(390)	422.16(388)

^aX-ray flux from 0.5–8.0 keV

light-curves do show evidence of moderate variability at the 3σ level. The longer *XMM-Newton* observation and superior high-energy response of EPIC-PN reveal strong emission lines of Fe XXV and Fe XXVI just below 7 keV. Weaker thermal emission lines are apparent between 1 and 3 keV (Figure 2).

The *XMM-Newton* spectrum was modeled by a Raymond-Smith thermal plasma model (Raymond & Smith 1977) with one or two temperature (kT) components absorbed by the intervening column density N_H . The metal abundance of the plasma was allowed to vary freely. The best-fit parameters, fluxes, 90% confidence intervals, and χ^2 values are given in Table 1. We confirm the low value of N_H and find a kT of $8.1^{+0.9}_{-0.8}$ keV given a single component. A second cooler component, $kT \sim 1$ keV, significantly improves the fit (F -test null-hypothesis probability $< 3.3 \times 10^{-6}$). However the flux in this component is $< 5\%$ of the flux in the hot component. Based on these spectra we conclude that the emission is dominated by an extremely hot thermal plasma with $kT = 10.4^{+2.3}_{-1.6}$ keV. At a distance to NGC 5281 of 1580 pc (Paunzen & Maitzen 2001) the measured N_H and fluxes imply a total X-ray luminosity of $6.2 \pm 0.1 \times 10^{32}$ erg s^{-1} . While a power-law model would of course fit the electron bremsstrahlung dominated spectrum, the feature just below 7 keV is well described by Fe XXV and Fe XXVI line emission in a thermal plasma at the temperature determined by the continuum.

4. DISCUSSION

Using the excellent spatial resolution of *Chandra* and optical images from *Magellan* we were able to identify the X-ray source 1WGA J1346.5–6255 thought to be a possible PWN at the center of SNR 309.2-0.6, as the O9.7e massive star HD 119682. The region itself is rather complex. The line of sight hosts an extended radio structure (Gaensler et al. 1998) at a distance of about 5 kpc which appears positionally coincident with a second possibly extended X-ray source from the *ROSAT* All Sky Survey (RASS) (Voges & et al. 1999) 1RXS J134651.1-624843. Follow-up *ASCA* observations detected strong emission-lines from the extended structure confirming its identification as a young ejecta-dominated SNR (Rakowski et al. 2001). As for the bright point source 1WGA J1346.5–6255, *ROSAT* contours already suggested the possibility of an association with the foreground cluster NGC 5281 and its third brightest star HD 119682 (Gaensler et al. 1998). However, its high X-ray brightness as well as a measured spectral index from




FIG. 2.— *XMM-Newton* spectra of 1WGA J1346.5–6255, from top to bottom, the single event PN spectrum ($\times 10$), the multiple event PN spectrum, MOS 1 ($\times 0.2$) and MOS2 ($\times 0.05$). The model shown includes two temperature components. The contribution of the second cooler component is illustrated for the MOS2 spectrum (marked “low kT ”). Fit residuals ($\delta\chi^2$) for each instrument are shown below the spectra.

the *ASCA* observation were equally consistent with a pulsar or synchrotron nebula interpretation (Rakowski et al. 2001). The low absorbing column density favored an origin in NGC 5281.

The better than $1''$ agreement between the *Chandra* source position and the optical position of HD 119682 dispels any doubts about the identification of 1WGA J1346.5–6255 with HD 119682. Both the optical and X-ray properties of HD 119682 are reminiscent of γ Cas where a recent X-ray gratings study concluded that the Be-star itself is the likely source of X-ray emission (Smith et al. 2004). First, HD 119682 has been classified as a very late O or very early B emission line star, similar in type to γ Cas (B0.5e). Second, the likeness between the X-ray properties of HD 119682 and γ Cas is striking. Our spectral analysis produced a two component fit that features a 1 keV soft component and a dominant hard component of about 10 keV. Significant Fe XXV and Fe XXVI line emission highlights the hot thermal origin of the spectrum in HD 119682. Likewise γ Cas exhibits a very hot emission component up to $kT = 12$ keV along with some levels of soft emission. In both stars the hard emission dominates, with the soft component contributing only 5% of

the flux in HD 119682 and 14% in γ Cas (Smith et al. 2004), which to our knowledge is unique in stellar systems. Furthermore, the measured luminosities are quite similar, $L_X = 6.2 \times 10^{32} \text{ erg s}^{-1}$, $\log(L_X/L_{bol}) = -5.4$ for HD 119682 compared to $L_X = 5.1 \times 10^{32} \text{ erg s}^{-1}$ (Kubo et al. 1998), $\log(L_X/L_{bol}) = -5.4$ for γ Cas using L_{bol} from Hamann & Persson (1992). That the emission from HD 119682 is analogous to γ Cas, seems by far the most likely possibility.

By contrast, the colliding wind scenario would have to involve another similarly massive companion with extremely strong winds, no evidence for which is seen optically in imaging or spectroscopy. Furthermore the spectral type of HD 119682 is O9.7e or later. For such late type stars the energetics for colliding winds as the origin of the hard X-ray emission are even more difficult to justify. A recent study by Schulz et al. (2006) involving the massive triple θ^2 Ori A, which contains an O-star of about similar type, showed that even for the existence a close massive companion, the wind densities as well as the involved kinetics do not favor a colliding wind scenario. HD 119682 should be even less favorable as optical spectra do not indicate a massive companion.

The possibility that the origin of the high-energy emission from HD 119682 is a magnetically confined wind seems a viable alternative. However, conditions that lead to a confinement scenario, as modeled for the case of θ^1 Ori C (Gagné et al. 2005), seem a bit unrealistic here given the extremely high plasma temperatures of HD 119682. Furthermore most magnetic activity of such kind has been observed in much younger OB stars (Schulz et al. 2003). Quite recently it was concluded that the field in young τ Sco, another young early type star that shows hard X-ray emission (Cohen et al.

2003), is likely of fossil origin rather than some intrinsic dynamo activity (Donati & et al. 2006). The answer thus has to be more complex.

The remaining possibility points to the quite enigmatic case of γ Cas. For the moment, though, with only a CCD-resolution spectrum we cannot go much beyond a discussion about the similarities between γ Cas and HD 119682. However, the case may bear implications with respect to the nature of the X-rays in both sources. HD 119682 is quite well identified and neither optical spectra (Levenhagen & Leister 2004) nor our high spatial resolution optical imaging yet show any evidence for a companion. Even though the high resolution spectra of γ Cas provided additional evidence that the observed X-ray emission is likely not due to a possible white dwarf companion, that hypothesis could not be entirely ruled out because of some resemblance between the X-ray emission from some cataclysmic variables and the HETG spectra presented by Smith et al. (2004). More recently Smith & Balona (2006) reported on another Be-star (B1e), HD 110432 (BZ Crucis), which shares similarly peculiar X-ray and optical characteristics with γ Cas. The case of HD 119682 thus seems to be another candidate of “ γ Cas analogs” with the strong notion that the hard X-ray emission emanates from a single OB emission line star.

C.E.R. was supported during this work by NASA Grant NAG5-9281. S.J.W. acknowledges support from CXC guest investigator grant GO5-6002A (ANCHORS) and NASA contract NAS8 39073 (CXC).

Facilities: Magellan:Clay (MagIC), CXO (ACIS), XMM (EPIC)

REFERENCES

- Berghoefer, T. W. & Schmitt, J. H. M. M. 1994, *A&A*, 292, L5
 Berghoefer, T. W., Schmitt, J. H. M. M., & Cassinelli, J. P. 1996, *A&AS*, 118, 481
 Cassinelli, J. P., Cohen, D. H., Macfarlane, J. J., Sanders, W. T., & Welsh, B. Y. 1994, *ApJ*, 421, 705
 Cassinelli, J. P., Miller, N. A., Waldron, W. L., MacFarlane, J. J., & Cohen, D. H. 2001, *ApJ*, 554, L55
 Chlebowski, T., Harnden, F. R., & Sciortino, S. 1989, *ApJ*, 341, 427
 Cohen, D. H., de Messières, G. E., MacFarlane, J. J., Miller, N. A., Cassinelli, J. P., Owocki, S. P., & Liedahl, D. A. 2003, *ApJ*, 586, 495
 Corcoran, M. F. 2005, *AJ*, 129, 2018
 Donati, J.-F. & et al. 2006, *MNRAS*, submitted
 Feldmeier, A. 1995, *A&A*, 299, 523
 Feldmeier, A. & Shlosman, I. 2002, *ApJ*, 564, 385
 Gaensler, B. M., Green, A. J., & Manchester, R. N. 1998, *MNRAS*, 299, 812
 Gagné, M., Oksala, M. E., Cohen, D. H., Tonnesen, S. K., ud-Doula, A., Owocki, S. P., Townsend, R. H. D., & MacFarlane, J. J. 2005, *ApJ*, 628, 986
 Hamann, F. & Persson, S. E. 1992, *ApJ*, 394, 628
 Harnden, F. R. & et al. 1979, *ApJ*, 234, L51
 Howk, J. C., Cassinelli, J. P., Bjorkman, J. E., & Lamers, H. J. G. L. M. 2000, *ApJ*, 534, 348
 Kahn, S. M., Leutenegger, M. A., Cottam, J., Rauw, G., Vreux, J.-M., den Boggen, A. J. F., Mewe, R., & Güdel, M. 2001, *A&A*, 365, L312
 Kubo, S., Murakami, T., Ishida, M., & Corbet, R. H. D. 1998, *PASJ*, 50, 417
 Levenhagen, R. S. & Leister, N. V. 2004, *AJ*, 127, 1176
 —. 2006, *MNRAS*, in press, astro-ph/0606149
 Lucy, L. B. 1982, *ApJ*, 255, 286
 Lucy, L. B. & White, R. L. 1980, *ApJ*, 241, 300
 Martins, F., Schaerer, D., & Hillier, D. J. 2002, *A&A*, 382, 999
 Miller, N. A., Cassinelli, J. P., Waldron, W. L., MacFarlane, J. J., & Cohen, D. H. 2002, *ApJ*, 577, 951
 Mink, D. J. 2002, in *ASP Conf. Ser. 281: Astronomical Data Analysis Software and Systems XI*, ed. D. A. Bohlender, D. Durand, & T. H. Handley, 169
 Owens, A., Oosterbroek, T., Parmar, A. N., Schulz, R., Stüwe, J. A., & Haberl, F. 1999, *A&A*, 348, 170
 Owocki, S. P., Castor, J. I., & Rybicki, G. B. 1988, *ApJ*, 335, 914
 Pallavicini, R., Golub, L., Rosner, R., Vaiana, G. S., Ayres, T., & Linsky, J. L. 1981, *ApJ*, 248, 279
 Paunzen, E. & Maitzen, H. M. 2001, *A&A*, 373, 153
 Pollock, A. M. T., Corcoran, M. F., Stevens, I. R., & Williams, P. M. 2005, *ApJ*, 629, 482
 Rakowski, C. E., Hughes, J. P., & Slane, P. 2001, *ApJ*, 548, 258
 Raymond, J. C. & Smith, B. W. 1977, *ApJS*, 35, 419
 Rho, J., Ramírez, S. V., Corcoran, M. F., Hamaguchi, K., & Lefloch, B. 2004, *ApJ*, 607, 904
 Sana, H., Stevens, I. R., Gosset, E., Rauw, G., & Vreux, J.-M. 2004, *MNRAS*, 350, 809
 Sanner, J., Brunzendorf, J., Will, J.-M., & Geffert, M. 2001, *A&A*, 369, 511
 Schulz, N. S., Canizares, C., Huenemoerder, D., & Tibbets, K. 2003, *ApJ*, 595, 365
 Schulz, N. S., Canizares, C. R., Huenemoerder, D., & Lee, J. C. 2000, *ApJ*, 545, L135
 Schulz, N. S., Testa, P., Huenemoerder, D., Ishibashi, B., & Canizares, C. R. 2006, *ApJ*, submitted
 Seward, F. D., Forman, W. R., Giacconi, R., Griffiths, R. E., Harnden, F. R., Jones, C., & Pye, J. P. 1979, *ApJ*, 234, L55
 Smith, M. A. & Balona, L. 2006, *ApJ*, 640, 491
 Smith, M. A., Cohen, D. H., Gu, M. F., Robinson, R. D., Evans, N. R., & Schran, P. G. 2004, *ApJ*, 600, 972

- Spitzbart, B. D., Wolk, S. J., & Bizunok, N. S. 2005, in *Protostars and Planets V*, 8518
- Townsley, L. K., Feigelson, E. D., Montmerle, T., Broos, P. S., Chu, Y.-H., & Garmire, G. P. 2003, *ApJ*, 593, 874
- Voges, W. & et al. 1999, *A&A*, 349, 389
- Waldron, W. L. & Cassinelli, J. P. 2001, *ApJ*, 548, L45
- White, N. E., Giommi, P., & Angelini, L. 1994a, *BAAS*, 26, 1372
- . 1994b, *IAU Circ.*, 6100, 1
- Wolk, S. J., Bourke, T. L., Smith, R. K., Spitzbart, B., & Alves, J. 2002, *ApJ*, 580, L161